



The 4 per 1000 goal and soil carbon storage under agroforestry and conservation agriculture systems in sub-Saharan Africa



Marc Corbeels^{a,b,*}, Rémi Cardinael^{a,c}, Krishna Naudin^{a,d}, Hervé Guibert^a, Emmanuel Torquebiau^a

^a CIRAD, UR Agro-Ecology and Sustainable Intensification of Annual Crops (AIDA), University of Montpellier, 34398 Montpellier, France

^b CIMMYT, Sustainable Intensification Program (SIP), P.O. Box 1041–00621, Gigiri, Nairobi, Kenya

^c University of Zimbabwe, Crop Science Department, Box MP167, Mt. Pleasant, Harare, Zimbabwe

^d Embrapa Cerrados, Rodovia BR-020, Km 18, P.O. Box 08223, CEP: 73310-970, Planaltina, DF, Brazil

ARTICLE INFO

Keywords:

Alley cropping
Carbon sequestration
Climate change mitigation
Crop residues
Crop rotation
Fallow
No-tillage
Parklands
Intercropping

ABSTRACT

The 4‰ initiative launched by the French government at COP21 in Paris in December 2015 aspires to increase global soil organic carbon (SOC) stocks at a rate of 0.4% per year. We conducted a systematic literature review on SOC storage under agroforestry and conservation agriculture systems in sub-Saharan Africa, where we reported 66 and 33 cases for both systems respectively. The results showed that SOC storage rates were significantly higher than 4‰ yr⁻¹ in fallows and in multistrata agroforestry systems ($P = 0.0001$ and 0.0178 , respectively), but not in alley cropping and parklands systems. For conservation agriculture, SOC storage rates were only significantly higher than 4‰ yr⁻¹ ($P = 0.0438$) when all three principles were applied, i.e. no- or minimum tillage combined with crop residue retention and intercropping or rotation. The data showed very large variability in SOC storage rates as the result of various factors, including previous land-use history, experimental set up and approach used to determine SOC storage (diachronic versus synchronic approach), soil type, depth of soil sampling, type of crops and management, and duration of the experiment. SOC storage rates significantly decreased with time in the agroforestry systems ($P = 0.0328$). However, we were unable to find significant relationships with initial SOC stocks or tree density. Given the limited published data and the high variability in results, no significant relationships between SOC storage rates and site variables were found for conservation agriculture. We argue that there is a potential for SOC storage in agricultural soils of sub-Saharan Africa, as illustrated by SOC gaps observed on smallholder farms. Low SOC levels are, however, to a great extent the result of limited resources of most smallholder farmers. Practices such as agroforestry and conservation agriculture can restore SOC in these soils, but the 4‰ initiative has to be implemented on the grounds of the positive impact on crop productivity rather than on climate change mitigation. The efficiency in doing so will depend on the specific situations and will need economic support to smallholder farmers, including the promotion of good markets for sale of extra produce and for input supply, effective private support and policy, such as credit schemes and subsidies for inputs, and efficient extension services which incentivize farmers to invest in new technologies.

1. Introduction

The Paris Climate Agreement of the United Nations Framework Convention on Climate Change (COP21, November 30 to December 11, 2015) commits to keeping the increase in global average temperature to well below 2 °C above pre-industrial levels and sets a more aspirational goal to limit warming to 1.5 °C. To achieve this, major greenhouse gas (GHG) reductions are needed together with technologies that result in net removal of CO₂ from the atmosphere (Meinshausen et al., 2009; Field and Mach, 2017). In this context, the “4 per 1000” initiative – “Soils for Food Security and Climate” – (<http://4p1000.org>) was

launched by the French Minister of Agriculture at the COP21, with the objective to increase soil organic carbon (SOC) stocks globally by 0.4 percent per year as a way to mitigate climate change and improve food security (Lal, 2016; Chabbi et al., 2017; Dignac et al., 2017; Minasny et al., 2017).

Soils store two to three times more carbon than the atmosphere, and the amount of organic carbon stored in soils is estimated at 2400 Gt C to 2 m depth (Batjes, 1996). A slight increase or decrease of this SOC stock has therefore a large impact on atmospheric CO₂ concentration (Schlesinger, 1977). Besides, the annual GHG emission from fossil carbon combustion is estimated at 9.3 Gt C (Le Quééré et al., 2016). The

* Corresponding author at: CIMMYT c/o ICRAF House, P.O. Box 1041-0062, Village Market, Gigiri, Nairobi, Kenya.
E-mail address: corbeels@cirad.fr (M. Corbeels).

4‰ figure simply originated from the ratio of the global anthropogenic carbon emissions (9.3 Gt C yr^{-1}) to the global SOC stock (2400 Gt C) (Minasny et al., 2017), and thus implies that increasing global SOC stocks by 4‰ per year would completely offset the anthropogenic CO_2 emissions. However, increasing SOC up to 2 m through changes in land management practices is rather impossible, because the management practices are known to principally affect the topsoil SOC content (VandenBygaert et al., 2018). Another interpretation of the 4‰ figure considers therefore the global SOC stock to 40 cm soil depth only, arguing that a more correct scientific rationale behind the 4‰ initiative is to counteract the annual rise in atmospheric CO_2 (de Vries, 2018). A 4‰ increase of the global 0–40 cm SOC stock, which is estimated at 860 Gt , would result in an annual sink of 3.4 Gt C yr^{-1} . Combining this with stopping deforestation (that causes an estimated emission of 0.9 Gt C yr^{-1} , Le Quéré et al., 2015) would lead to a net CO_2 removal of 4.3 Gt C yr^{-1} from the atmosphere, close to the present annual rise in atmospheric CO_2 , being estimated at 4.5 Gt C (Le Quéré et al., 2016).

Soil organic carbon stocks of Africa are currently estimated between 133 and 184 Gt for the 0–100 cm soil layer (about 10% of the global SOC stock), with large variations depending on the ecoregion (Henry et al., 2009; Scharlemann et al., 2014). Due to an increasing demographic pressure and associated food demand, many native grassland and forest areas have been converted to croplands, and soils have experienced a massive decline in SOC, mainly as a result of low carbon inputs into soil, and because of topsoil erosion (Agbenin and Goladi, 1997; Solomon et al., 2002; Lemenih et al., 2005; Zingore et al., 2005; Kintché et al., 2010; Moebius-Clune et al., 2011). For example, from a review of published research it was estimated that SOC in croplands in southern Africa has declined by 25 to 53% (Swanepoel et al., 2016). Other studies (e.g. in Western Kenya) found decreases of up to 85% of the original SOC stocks under forest (Solomon et al., 2007). These soils, far from their saturation level (Six et al., 2002; Stewart et al., 2007), are now seen as important potential carbon sinks. The concept of SOC saturation suggests that the quantity of stable SOC in a given soil is limited since SOC stabilization, i.e. SOC protection from decomposers, is to a great extent determined by the amount of fine soil particles (typically $< 20 \mu\text{m}$) (Hassink, 1997; Six et al., 2002). The difference between the theoretical SOC saturation value and measured SOC of the fine fraction corresponds to the soil saturation deficit and may represent the potential for SOC storage in a stable form (Castellano et al., 2015), but this concept is now being challenged (Barré et al., 2017).

Soil organic carbon increase can be achieved through the implementation of restorative land use and recommended management practices on agricultural soils (Lal, 2004). Restoring SOC levels is often seen as a win-win strategy to mitigate climate change while improving soil quality, potentially leading to higher crop yields (Lal, 2006). Important effects of increased SOC levels are increased supply of nutrients (Vanlauwe et al., 2005; Bationo et al., 2007), and better soil structure resulting in increased infiltration of water and reduced soil crusting, compaction and erosion (Blanco-Canqui and Lal, 2007; Nyamadzawo et al., 2007). Higher SOC levels also result in more efficient use of the agricultural inputs in cropping systems (Tittonell et al., 2008; Zingore et al., 2011). This in turn may lead to improved climate change adaptation, increased food security and better livelihoods (Lal, 2004).

Several management practices are recognized to improve SOC contents in croplands, such as organic amendments, cover crops, improved crop rotations with or without temporal grasslands or legumes, biochar, agroforestry, or conservation agriculture (Vågen et al., 2005; Lorenz and Lal, 2014; Paustian et al., 2016).

This study focuses on the impact of agroforestry and conservation agriculture on SOC storage. Both practices can increase SOC through increased carbon inputs from higher biomass productivity within a field and do not lead to carbon leakage from one field to another as often the case with recycling of exogenous organic matters (Powlson et al., 2011).

The first aim of this study is to develop evidence for increased SOC

stocks under agroforestry and conservation agriculture systems in sub-Saharan Africa based on a systematic review of available data in the literature. We present the results of the review against the 4‰ objective and discuss the extent to which agroforestry and conservation agriculture can enhance SOC storage and contribute to climate change mitigation. Second, we highlight the opportunities and challenges that lay ahead for SOC storage in the context of African smallholder farming systems by virtue of findings from studies on soil fertility gradients on smallholder farms. Finally, we discuss the challenges with the widespread implementation and monitoring of the 4‰ initiative in the context of African farming systems, leading to concluding commentaries on the overall findings of our study.

2. Materials and methods

2.1. Literature search

Using the online databases ISI Web of Science (Thomson Reuters, New York, NY, USA) and Scopus (Elsevier, Amsterdam, the Netherlands), a comprehensive search was conducted in May 2017 for peer-reviewed publications that reported SOC storage under agroforestry and conservation agriculture systems in sub-Saharan Africa (excluding South Africa). The literature search was conducted using the terms described in Table S1 (Supplementary Materials). Agroforestry is defined as the intentional, integrated combination of trees (including shrubs) and agricultural crops simultaneously or sequentially grown on the same piece of land (Nair, 1993; Torquebiau, 2000). Simultaneous systems include scattered trees in cropland, often referred to as ‘parklands’, alley cropping, intercropping and multistrata agroforestry. Sequential systems include improved fallows, relay cropping with trees and rotational woodlot systems where a piece of land is deliberately planted with fast-growing, often nitrogen-fixing trees (Nair, 1985). Conservation agriculture is defined according to FAO (2015) as a cropping system characterized by three principles, namely 1) direct seeding of crops with minimal soil disturbance; 2) retention of crop residues as mulch on the soil surface, and 3) the use of crop rotations and/or intercropping.

Retrieved publications were screened and had to meet the following defined basic selection criteria. First, studies are field experiments conducted in sub-Saharan Africa in which SOC measurements were done on a control treatment (i.e. a cropland without an agroforestry or conservation agriculture intervention) and on a pairwise agroforestry treatment or a (partial) conservation agriculture treatment where at least the effect of no- or minimum tillage was tested (‘space for time’ or synchronic approach), or on an agroforestry or conservation agriculture treatment with repeated soil sampling in time (diachronic approach). Results from chronosequence studies and studies using forests or natural ecosystems as a control were excluded. Second, the same crop was grown in both treatment and control. Third, studies report SOC stocks, which is the quantity of carbon per unit area expressed in kg C m^{-2} or kg C ha^{-1} , or SOC stocks could be calculated from the published SOC concentration expressed as proportion of the total soil mass, and soil bulk density. Studies that reported only SOC concentrations without soil bulk density were excluded. In some cases when published data were incomplete to calculate SOC stocks, the authors of the studies were contacted for additional information. In this way, SOC stocks were retrieved from the original datasets provided by the authors for the following studies: Baumert et al. (2016), Bright et al. (2017), Paul et al. (2015) and Rimhanen et al. (2016). Fourth, the treatment duration is reported in the paper.

For each study, we calculated the absolute annual rate of change in SOC stock ($\text{kg C ha}^{-1} \text{ yr}^{-1}$) and the relative rate (i.e. to the initial or control value of the SOC stock, $\% \text{ yr}^{-1}$) to the maximum soil depth specified from the initial and final measurements (diachronic approach) or from the SOC stocks under treatment and control (synchronic approach), and the duration of the treatment. When SOC storage rates




Fig. 1. Number of observations for agroforestry and conservation agriculture as a function of treatment age and soil sampling depth.

could be calculated with both the diachronic and synchronic approach, we used the diachronic rate in the further analysis of data because confounding effects of spatial heterogeneity with the synchronic approach may lead to less accurate results compared to the diachronic approach (Costa et al., 2013). Four studies on agroforestry systems reported SOC contents for soil depths well below 40 cm, i.e. up to 100 or 200 cm soil depth. To account for these potential soil depth effects, we also calculated and analyzed SOC storage rates to a maximum soil depth of 40 cm. If SOC contents to 40 cm depth (or less) were reported in the above four studies, they were included in this additional analysis. If a site was sampled more than once at different points in time, we used the results of the first and last reported samplings, so to obtain the largest sampling interval. Data from the same study but reported in more than one publication were not repeated to avoid double counting; the publication with the most complete dataset was used. When the study investigated multiple agroforestry or conservation agriculture treatments, each treatment was considered as a single observation in the analysis. When multiple controls were reported, the most representative practice for the study area was used as control, and when more than one form of tillage was assessed in a study on conservation agriculture, we selected the tillage treatment representing the greatest soil disturbance as control. When a paper presented results from single control and treatment plots (no replicates), the study was not included in the analysis.

The literature search returned 183 publications for agroforestry and 157 for conservation agriculture. With the above selection criteria, 15 and 20 publications were selected in the final dataset for agroforestry and conservation agriculture, respectively. The complete reference list and details of selected publications are given as Supplementary Material (Tables S2, S3).

2.2. Statistical analyses

All the statistical analyses were performed using R software version 3.1.1 (R Development Core Team, 2013), at a significance level of $P \leq 0.05$. Reported data concerning SOC storage rates for the different agroforestry and conservation agriculture systems presented heterogeneous variances, preventing from using classical statistical tests (Scheffé, 1959; Littell et al., 2006). We therefore used a weighted least squares regression, using the *gls* function from the *nlme* package (Galecki and Burzykowski, 2013). Each observation per subsystem *i* (alley cropping, fallows, multistrata system, parkland; no- or minimum tillage, no- or minimum tillage + residues, no- or minimum

tillage + residues + rotation/intercropping) was weighted by $\frac{1}{V_i}$, where V_i was the variance of a given subsystem. Weights were normalized so that their sum equaled the total number of observations ($n = 66$ for the agroforestry, $n = 33$ for conservation agriculture). The *lsmmeans* function (Lenth, 2016) was then used to test whether SOC storage rates for each subsystem were significantly different from $0\% \text{ yr}^{-1}$ and from $4\% \text{ yr}^{-1}$. To compare the different subsystems, pairwise comparisons were also performed with the *lsmmeans* function, using the Tukey's method.

3. Results

3.1. Agroforestry

The total of 15 retrieved publications from peer-reviewed literature resulted in 68 observations with absolute changes in SOC stocks reported. For two observations (Verchot et al., 2011) the relative change of SOC stock could not be calculated since the control SOC levels were not reported (Table S2). The final review deals with the 66 remaining observations from 12 countries. There were 19 observations from alley cropping and eight from intercropping systems which we regrouped in our analysis. Nineteen observations came from improved fallows, one from natural fallow and five from rotational woodlots which we combined as 'fallows'. Finally, eight observations were from multistrata systems and six from parklands. Only two publications (Raddad et al., 2006; Bright et al., 2017) presented SOC stocks measured at different time periods (diachronic approach); all other publications reported data from paired-site experiments. Since the synchronic approach in the study from Bright et al. (2017) allowed to calculate SOC storage rates that covered a longer time span (13 years) than the diachronic approach (4 years), we considered the synchronic rates for this study in the further analysis. Maximum soil sampling depth ranged from 12 to 200 cm (Fig. 1). A majority of the data (62%) covered only the 20 first centimeters of soil, while 20% of the data reported a maximum sampling depth of 30 to 50 cm. Only 14% of the observations were obtained by sampling at least down to 1 m depth. The time since the introduction of the agroforestry system varied from 2 to 50 years, with 70% of the observations that had trees planted less than 10 years before the final soil sampling (Fig. 1).

Overall, the absolute rate of SOC change varied from -9.6 ($0-40$ cm) to $14 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ ($0-100$ cm) (Table S2). The largest rate of SOC increase was from a study with 11 years of continuous alley cropping with *Maesopsis eminii* Engl. on a sandy loam in Uganda (Tumwebaze et al., 2011). The largest decrease reported was on a




Fig. 2. Cumulative distribution function of SOC storage rates in agroforestry and in conservation agriculture systems.

loamy sand site in Burkina Faso under a 2-year alley cropping system with *Jatropha curcas* L. (Baumert et al., 2016).

Relative rates of SOC change varied widely from -159 to $+194\text{‰ yr}^{-1}$ (Fig. 2, values considering maximum reported soil depths). Eighty percent of the observations had annual relative SOC storage rates higher than 4‰ yr^{-1} . Fifty percent of the observations had SOC storage rates of 34‰ yr^{-1} or higher.

The alley cropping, fallows and multistrata systems showed similar median rates of, respectively, 37, 37 and 45‰ yr^{-1} . Alley cropping systems showed the highest variability, with storage rates ranging from -159 to 149‰ yr^{-1} (Fig. 3). Most negative rates of SOC change were from paired-site comparisons between sorghum cropped alone and alley

cropped with *Jatropha curcas* L. at seven sites in Burkina Faso with the age of the alley cropping system varying between two and four years (Baumert et al., 2016). Four observations of this same study showed, however, positive rates of SOC storage of between 15 and 67‰ yr^{-1} . Two negative rates were also observed in multistrata systems in Ethiopia (Rimhanen et al., 2016). The highest rates of SOC storage under alley cropping ($72\text{--}149\text{‰ yr}^{-1}$) were from paired-site comparisons where the soils were sampled to, respectively, 1 m and 2 m depth, i.e. the four observations reported by Tumwebaze et al. (2011) on alley cropping with four different tree species, and the two observations reported by Makumba et al. (2007) on alley cropping with *Gliricidia sepium*. We retrieved only one publication on multistrata systems




Fig. 3. Soil organic carbon storage rates for different agroforestry and conservation agriculture systems. Upper and lower edges of boxes indicate 75th and 25th percentiles, horizontal lines within boxes indicate median, whiskers below and above the boxes indicate the 10th and 90th percentiles, and crosses indicate mean SOC storage rate per type of subsystem. Outliers are plotted as individual points. For each system, different letters represent significant differences ($P \leq 0.05$) between subsystems. ** represents subsystems with SOC storage rates significantly ($P \leq 0.05$) higher than 0‰ yr^{-1} and 4‰ yr^{-1} .




Fig. 4. Comparison of SOC storage rates estimated by the diachronic or synchronic approach in conservation agriculture systems.

(Rimhanen et al., 2016) that reported eight paired-site comparisons (0–15 cm sampling depth) at different sites in Ethiopia. Soil organic carbon storage rates under these systems ranged widely from -5 to $80\% \text{ yr}^{-1}$. For parkland systems, there were six observations from two publications (Abaker et al., 2016, 5 observations; Gelaw et al., 2014, 1 observation). The SOC storage rates varied from -3 to $22\% \text{ yr}^{-1}$, mainly depending on the type of grass species in the treatments.

Soil organic carbon storage rates were significantly higher than $4\% \text{ yr}^{-1}$ (and $0\% \text{ yr}^{-1}$) in the fallows and in the multistrata systems ($P = 0.0001$ and 0.0178 , respectively) (Fig. 3, values considering maximum reported soil depths). Rates in the fallows were significantly higher than in the parklands ($P = 0.0003$). When we considered reported SOC contents to a maximum soil depth of 40 cm only, SOC rates in the fallows became significantly higher than in the alley cropping systems ($P = 0.0169$) (Fig. S1, Supplementary Materials). For the parkland systems, we were left with only one observation. Rates in the fallows and in the multistrata systems remained significantly higher than $4\% \text{ yr}^{-1}$ (and $0\% \text{ yr}^{-1}$) ($P = 0.0001$ and 0.0144 , respectively).

3.2. Conservation agriculture

We retrieved 20 publications that resulted in a total of 33 observations, comprising 21 sites in seven countries (Table S3). Maize was by far the most studied crop species (27 observations); the remaining studies related to six other crop species. In several studies, only one or two of the conservation agriculture principles were applied: five observations were from treatments with no-tillage alone and three observations from no- or minimum tillage with intercropping and/or rotations, but without crop residue mulching. They were grouped in the analysis (Fig. 3). Ten observations were from treatments with no- or minimum tillage with crop residue mulching, but without rotation or intercropping, and 14 observations referred to treatments with all three principles of conservation agriculture applied. In one study (Ojeniyi and Adekayode, 1999), it was not clear whether crop residues were left on the soil or not. The one observation from this study was therefore not used when differentiating between the varying degrees of application of the conservation agriculture principles (Fig. 3). Most conservation agriculture treatments in the dataset involved no-tillage, i.e. manual or mechanized direct seeding (26 observations). The other seven observations were minimum tillage treatments, i.e. ripping (5

observations) or surface scraping of the soil (2 observations). The tillage type to which the conservation agriculture treatment was compared was mainly ploughing (11 observations with ox-ploughing and 9 observations with tractor ploughing). Hoeing and ridging represented respectively eight and five observations. Fourteen studies measured SOC storage through repeated sampling in time of the conservation agriculture treatment, but at the same time these studies were also paired-site comparisons between a conservation agriculture and conventional tillage treatment; the other six studies were only paired-site comparisons. Soil sampling depth ranged from 5 to 60 cm, but 91% of the observations had a maximum sampling depth of 30 cm (Fig. 1). The time since the introduction of the conservation agriculture treatment varied from 1 to 11 years, but 85% of the observations had a maximum treatment age of 5 years (Fig. 1).

Overall, the absolute SOC change varied from -10.2 (0–30 cm) to 6.86 (0–15 cm) $\text{Mg C ha}^{-1} \text{ yr}^{-1}$ for observations where SOC storage was calculated using the diachronic approach (Table S3). The largest decrease ($-10.2 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$) reported was from a 2-year experiment on a site in south-eastern Nigeria where sweet potato was cultivated on a loam under no-tillage without residue retention (Anikwe and Ubochi, 2007). This large decrease can probably be explained by the fact that the site was slashed and cleared of fallow vegetation, and thus had an initial high level of relatively labile SOC that decomposed rapidly following cultivation. The data also showed large variability with least significant difference ($P = 0.05$) values of up to 8.6 Mg C ha^{-1} . The largest rate of SOC increase ($6.86 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$, diachronic approach) was from a study in southern Nigeria on a sandy loam under 2-year no-tillage without residue retention (Agele et al., 2005), but this is probably an overestimate due to the small soil sampling depth (0–15 cm); the practice of no-tillage results in stratification of SOC concentration with depth, with surface soil layers having considerably higher SOC concentration than the layers below (Baker et al., 2007).

Relative SOC change varied from -142 to $450\% \text{ yr}^{-1}$ (Fig. 2). SOC change under conservation agriculture was negative for nine observations. Sixty-four percent of the studies had annual relative SOC storage rates that were higher than $4\% \text{ yr}^{-1}$. Fifty percent of the observations had relative rates that were larger than $34\% \text{ yr}^{-1}$. Treatments with no-tillage without residue retention showed the smallest median SOC storage rate, $-12\% \text{ yr}^{-1}$ (Fig. 3), and the highest variability, ranging from -142 to $450\% \text{ yr}^{-1}$. The reported negative rates of SOC change were from repeated sampling on sites that were cleared from native vegetation. They became however positive when calculated on the basis of the comparison with the conventional tillage treatment (Fig. 4). Rates under no- or minimum tillage with residue retention also varied widely between -98 and $448\% \text{ yr}^{-1}$. Again, negative rates were reported from two experiments in Nigeria where the conservation agriculture treatment was installed following clearing of native vegetation (Agbede, 2008; Lal, 1998). The highest rate ($448\% \text{ yr}^{-1}$) was reported in a paired-site study in Malawi (Mloza-Banda et al., 2014) in which manual direct seeding with retention of crop residues was compared with ridging without crop residues on three different fields after four years. The value is an outlier and probably an overestimate. Apart from the methodological problems with making comparisons of SOC stocks between flat planting and ridge and furrow systems, the data were from three paired-site comparisons on different farmer fields where great variability between fields is to be expected. Annual SOC rates from treatments with all three principles of conservation agriculture varied somewhat less, between -96 and $176\% \text{ yr}^{-1}$ (Fig. 3). The lowest rate ($-96\% \text{ yr}^{-1}$) was reported in a 2-year study in Zimbabwe by Nyamadzawo et al. (2008) in which SOC storage was measured on a sandy clay loam (0–20 cm) under direct seeding of maize, following a fallow period with *Sesbania sesban*. The highest value of $176\% \text{ yr}^{-1}$ was from a sandy loam site in Western Kenya (Ngome et al., 2011) where the soil (0–15 cm) was sampled after 1 year in a no-tillage treatment with maize grown on a permanent living cover of *Arachis pintoii*. Two other observations from the same study with the same

conservation agriculture treatment at two different sites gave also relatively high rates of $102\% \text{ yr}^{-1}$ (on a clay) and $77\% \text{ yr}^{-1}$ (on a clay loam). The values were however derived from small absolute changes in measured SOC concentrations (0.19, 0.16 and 0.15%, respectively). The increase is certainly also the result of SOC redistribution nearer to the soil surface.

Only SOC storage rates in the system with the full set of conservation agriculture principles, i.e. no- or minimum tillage, residue retention and intercropping or rotation were significantly higher than $0\% \text{ yr}^{-1}$ ($P = 0.0244$) and $4\% \text{ yr}^{-1}$ ($P = 0.0438$). Soil organic carbon storage rates were not significantly different between the different degrees of application of the conservation agriculture principles (Fig. 3).

The comparison of studies for which SOC storage rates could be calculated both using the diachronic and the synchronic approach showed that the synchronic rates were in most cases higher than $0\% \text{ yr}^{-1}$ (two exceptions). Synchronic rates were much larger than diachronic rates that showed negative values (Fig. 4). These large deviations were from conservation agriculture experiments that were installed following clearance of native vegetation or fallow. In these cases, initial SOC was relatively high, and probably labile provoking net negative changes in SOC, both for the treatment and control.

4. Discussion

4.1. SOC storage potentials

Agricultural soils in sub-Saharan Africa have a strong carbon sink capacity as a result of previous SOC depletion (Lal, 2016). Theoretically, SOC stocks can be restored through soil and crop management practices that have a positive carbon budget (i.e. with carbon inputs from biomass exceeding carbon outputs through decomposition and erosion (Stockmann et al., 2013; Dignac et al., 2017)). We examined the SOC storage potentials of two widely promoted management practices in sub-Saharan Africa, agroforestry and conservation agriculture. In the following sections we discuss the extent to which these practices can increase SOC stocks and contribute to climate change mitigation. Other practices that increase carbon inputs to soil are e.g. the addition of manure or other organic materials to fields. However, adding organic materials to soil does not necessarily lead to a net transfer of carbon from the atmosphere to the soil. Depending on the alternative end use of the materials, it can just constitute a transfer from one location to another, and thus without mitigating climate change (Powlson et al., 2011).

4.1.1. Agroforestry

Overall, our review showed that agroforestry has a potential to store SOC (Figs. 2 and 3), especially multistrata systems and fallows. The largest SOC storage rates ($> 190\% \text{ yr}^{-1}$) were obtained with rotational woodlots (Kimaro et al., 2011). These agroforestry systems conflict however with food security goals, since edible crops are not grown in these systems for prolonged periods of time. Parklands showed the lowest potential for storing SOC (Fig. 3), probably because of the low tree density per hectare (Takimoto et al., 2008). Besides improving carbon levels in soils, agroforestry systems store also a vast amount of carbon in the tree aboveground and belowground biomass, which considerably increases their interest in a perspective of climate change mitigation (Luedeling et al., 2011; Zomer et al., 2016; Feliciano et al., 2018).

Trees have an extensive root system that can grow deep into the soil (Cardinael et al., 2015b; Pierret et al., 2016) and root derived carbon is probably the most important source for SOC storage since it is more likely to be stabilized in the soil by physicochemical processes than shoot derived carbon (Balesdent and Balabane, 1996; Menichetti et al., 2015). The highest relative SOC storage rates ($72\text{--}149\% \text{ yr}^{-1}$) in our database were reported from experiments where soil was sampled to depths of 1 to 2 m, confirming the need to sample deep soil layers to study agroforestry systems (Haile et al., 2010; Cardinael et al., 2015a). Overall, trees enhance SOC storage because of higher biomass production and carbon inputs to soil (Peichl et al., 2006; Cardinael et al., 2018) caused by increased light, water and nutrient use efficiency at the plot scale compared to a monoculture of annual crops. Trees can also recover nutrients from deeper soil layers and thus improve soil nutrient availability and uptake (Nair et al., 1999). Likewise, agroforestry systems with N-fixing trees may cause higher biomass production and carbon inputs as a result of increased nitrogen availability (Sileshi et al., 2008). Effects of agroforestry systems on rates of SOC decomposition are far less understood (Lorenz and Lal, 2014). Changes in litter quality and in the soil microbial and faunal communities involved in SOC decomposition may play a role (Gnankamby et al., 2008; Mitchell et al., 2010).

Within a type of agroforestry system, published rates of SOC storage varied widely (Fig. 3, Table S2). The age of the agroforestry system, tree density, initial SOC stock and soil texture are amongst the most influencing factors that determine SOC storage rates (Laganière et al., 2010) and to a certain extent explain the variability in published data. Rates of SOC change are likely to be smaller for older agroforestry systems since they are expected to gradually decrease as a new SOC equilibrium is reached over time under the agroforestry system (Paustian et al., 1997). This also means that a strict interpretation of the 4% concept of




Fig. 5. Soil organic carbon storage rates as a function of initial SOC stocks, age since the establishment of the studied practice, and tree density.

increasing SOC stocks each year by a slightly larger absolute amount as SOC increases is not reasonable. A significant decrease of SOC storage rate with time was indeed found for agroforestry systems ($P = 0.0328$) (Fig. 5). This entails that the capacity for SOC storage is finite and net CO_2 removals under a new management system are thus of limited duration (Powelson et al., 2011; Paustian et al., 2016). On the other hand, we did not find significant relationships between SOC storage rate and initial SOC stock or tree density (Fig. 5), which is probably due to the fact that other factors, such as tree species, type of control treatment and soil and climatic conditions interfered. It is generally known that positive effects on SOC are more likely to occur in finer-textured soils than in sandy soils because of the physico-chemical stabilization of SOC by silt and clay particles (Torn et al., 1997), but we were not able to test for this effect due the lack of soil texture data in several of the retrieved publications.

In conclusion, our review suggests a high potential for SOC storage under newly established agroforestry systems in soils of sub-Saharan Africa, which represents a genuine option for climate change mitigation. Most of the published values were well above the $4\% \text{ yr}^{-1}$. The limited number of observations and the high variability in results ask for more research to better understand the positive effects of different agroforestry systems on SOC and how management could optimize these effects (Nair and Nair, 2014; Feliciano et al., 2018).

4.1.2. Conservation agriculture

Conservation agriculture has received much attention as a practice to increase SOC stocks and mitigate climate change (Hobbs and Govaerts, 2010; Lal, 2015). In general, SOC storage under conservation agriculture has been attributed to increased retention of crop residues compared to conventional cropping practices (Virto et al., 2012). For example, in a global meta-analysis, Luo et al. (2010) found that in most studies no-tillage alone did not increase SOC stocks; on the other hand, if the practice of no-tillage induced increased biomass production (through e.g. the cultivation of two crops per year instead of one), SOC stocks increased. These findings are corroborated by the results from our review and from a recent review by Powelson et al. (2016): the largest SOC storage rates are observed in no- or minimum tillage treatments with retention of crop residues and with the use of crop rotations or intercropping. We found that conservation agriculture produced SOC storage rates compatible with the “4 per 1000” initiative, only when all three principles are implemented (Fig. 3).

No-tillage systems with retention of crop residues are also effective in reducing carbon losses through soil erosion (Ranaivoson et al., 2017). Carbon lost on a field through erosion is, however, not necessarily converted to CO_2 as it may be transported and (permanently) stored in soil or sediments in another location (Wang et al., 2017).

No- or minimum tillage alone can lead to SOC storage if it decreases the rate of SOC decomposition. Six et al. (1999) showed that less disruption of macro-aggregates with no-tillage enhances the formation of micro-aggregates that are crucial in the stabilization (lower decomposition rate) and storage of SOC in the long term. The results from our review showed however that there was no positive effect of no-tillage on SOC storage (Fig. 3). Besides, it is important to note that conversion from conventional tillage to no-tillage alters the vertical distribution of SOC in the soil profile, resulting in increased SOC in the 0–20 cm soil layers and decreased soil carbon in the 20–40 cm soil layers (Baker et al., 2007; Angers and Eriksen-Hamel, 2008; Luo et al., 2010). This means that an observed increase of SOC in the topsoil layer under no-tillage does not necessarily represent a net increase in the overall SOC, and that studies on SOC storage should investigate at least the 0–40 cm soil layer and not only the topsoil.

Our review showed highly variable SOC storage under conservation agriculture, with some reported rates that were anomalously large or low (Figs. 2 and 3). A conclusion that was also made by Powelson et al. (2016) in their meta-analysis of SOC stock changes under conservation agriculture practices in the Indo-Gangetic Plains and SSA. We did not

find significant relationships between SOC storage rate and initial SOC stock or age of the conservation agriculture systems (Fig. 5). The effects of these factors may be confounded with the effects of several other factors that influence SOC storage rates. First, the agro-ecological conditions across the studies are highly heterogeneous. For example, diachronic studies that were established on a land that was cleared from natural vegetation will result in negative SOC storage rates even under conservation agriculture, whilst studies on degraded land may result in build-up of SOC. Second, various forms of conservation agriculture (and conventional tillage practices) were tested in the reported experiments. Third, several methodological approaches are used for measuring SOC storage under conservation agriculture, including diachronic versus paired-site approaches (Fig. 4); on station versus on farm experiments; and different depths of soil sampling (Fig. 1).

There was a major difference in SOC storage rates estimated by the diachronic versus the synchronic approach, the latter one usually leading to higher estimated rates (Fig. 4). This phenomenon can be explained by the land use history of the fields (Costa et al., 2013). The diachronic approach provides a measure of net SOC changes under a treatment with respect to an initial state. These changes are expected to be negative under any agricultural system that was recently installed on soils that were previously under natural ecosystems (e.g. Solomon et al., 2007). In other words, diachronic rates depend to a large extent on the initial levels (and quality) of SOC, and thus on the land-use history. On the other hand, the synchronic approach compares the difference in SOC stocks of a treatment with respect to a control. Spatial variability of soil properties can interfere in the synchronic approach where space is substituted for time (Costa et al., 2013). As such, positive synchronic SOC rates do not provide evidence of a net SOC storage or a true carbon sink that removes CO_2 from the atmosphere. This pleads for long-term diachronic experiments.

In conclusion, conservation agriculture with the three principles applied can increase SOC and offer a scope for climate change mitigation, if carbon inputs are increased through higher crop productivity (via intercropping and/or crop rotations). The impact will depend on the site conditions, type of cropping system, management and the historical land use. Importantly, conservation agriculture practices, and SOC-restoration practices in general, once established, need to be preserved over time. If the practice of CA that led to increased SOC stocks is reversed, the previously accumulated SOC will be lost as CO_2 , and usually more rapidly than it was accumulated (VandenBygaert and Kay, 2004).

4.2. Soil fertility gradients illustrate potential and challenges with SOC storage

The potential and challenges with SOC storage in the agriculture landscapes of sub-Saharan Africa are probably best illustrated with studies on soil fertility gradients on smallholder farms (Tittonell et al., 2013). Soil fertility gradients are a consequence of the differential allocation of nutrient and labor resources to the various fields of a farm. They are the result of the limited resources that smallholders possess and that they tend to concentrate in the fields close to their homestead. Over time, this leads to soil fertility gradients, including gradients of SOC. Soil fertility typically decreases within a farm at increasing distances from the homestead (Prudencio, 1993; Carter and Murwira, 1995; Tittonell et al., 2005; Zingore et al., 2007). Reinforcing the soil fertility gradients, farmers often prioritize nutrient and labor allocation to their most fertile fields, where crop responses to inputs are generally highest.

For example, a case study in Western Kenya (Tittonell et al., 2008) showed gradients of SOC levels across fields in farms at three different locations (Table 1). On average per location, SOC levels in the outfields were 61, 67 and 83% of those in homestead fields. The midfields had SOC levels that were 60, 73 and 88% of the SOC levels in the homestead fields. These SOC gaps between fields are mainly the results of the

Table 1
Gradients of soil organic carbon (SOC) levels (0–30 cm soil layer) and maize grain yields on farmer fields at three locations in Western Kenya (from [Tittonell et al. \(2008\)](#)).

Location/position within the farm	SOC (g kg ⁻¹)	Grain yield (kg ha ⁻¹) ^a
Aludeka		
Homestead field	10.9	1700
Midfield	6.6	1000
Outfield	6.7	700
Emuhaya		
Homestead field	17.4	2400
Midfield	12.8	2200
Outfield	11.7	1400
Shinyalu		
Homestead field	19.6	2600
Midfield	17.2	1700
Outfield	16.2	1400

^a Average maize grain yields during the 2002 long rains season.

higher use of manure or composted household waste on the homestead fields compared to the other fields. Besides, the higher soil fertility combined with better overall management (including the use of mineral fertilizers) of homestead fields gave rise to higher crop yields ([Table 1](#)), which in turn results in higher carbon inputs (via roots, and residues if left on the field). These findings suggest that there is a wide scope for storing SOC in the midfields and outfields; in the presented case study between 10–40% of the original SOC levels. On the other hand, SOC storage on those fields is not achieved given the limited production resources of the farmers. Several authors pointed out that SOC storage can only be achieved with increased nutrient (most importantly, nitrogen) availability in soils ([van Groenigen et al., 2006, 2017](#); [Kirkby et al., 2013, 2014](#); [de Vries, 2014](#); [Richardson et al., 2014](#)).

The above spatial effects of differential allocation of limited resources by farmers are also occurring at the larger scale of a village, as for example illustrated in a study in north-east Zimbabwe by [Zingore et al. \(2011\)](#). Wealthier farmers with cattle have usually fields with higher SOC levels (especially their homestead fields) as a result of application of manure compared to poorer farmers without cattle who do not have manure. Moreover, crop residues from fields of non-cattle farmers are often grazed by cattle of the wealthier farmers, reducing the carbon inputs in the soils of these fields. As a result, fields of non-cattle owners are SOC depleted. These fields represent an important carbon sink, if farmers' resources would allow to purchase and apply organic manures, alone or in combination with mineral fertilizer. Although the higher SOC storage in the fields of the wealthier cattle-owners are to a large extent the results of a transfer of organic material from the communal grazing areas and fields of non-cattle farmers to their own fields, and does not constitute a net transfer of carbon from atmosphere to soil, the higher SOC levels on fields of the wealthier farmers result in higher nutrient use efficiency and crop yields, thus providing more carbon inputs to the soil (via roots), and thus a net transfer of carbon from atmosphere to soil.

4.3. Implementation and adoption of SOC restoration practices

Strategies for widespread adoption of SOC-restoration practices are required for the 4‰ initiative to be effective in mitigating climate change. In sub-Saharan Africa, however, the challenge to implement these strategies for agricultural soils is enormous. Many farmers are extremely poor and live below the poverty line, often with the inability to invest in new practices or technologies. It should also be clear that for smallholders, climate change mitigation is no priority ([Luedeling and Neufeldt, 2012](#)). Farmers' priorities are to get immediate benefits from farming and to ensure their food security ([Jerneck and Olsson, 2014](#)). Therefore, the entry point for promoting SOC-restoration practices should be on the grounds of increasing crop productivity through

better soil quality, and not on the basis of climate change mitigation. The majority of the management practices that build SOC these practices can be promoted on a “no regrets” basis ([Smith et al., 2008](#)), i.e. they do provide other benefits even if the climate change mitigation potential is not realized. Especially in low-input smallholder farming systems, enhancing SOC storage positively impacts crop productivity (e.g. [Tittonell et al., 2008](#)), and is expected to contribute to enhanced resilience to climate change.

Despite these co-benefits, adoption of SOC-restoration practices, such as agroforestry and conservation agriculture, by smallholders is low in sub-Saharan Africa ([Giller et al., 2009](#); [Jerneck and Olsson, 2013](#)). There are several technical, infrastructural, socio-economic, or policy barriers to adoption. First of all, farm sizes are small (often less than 2 ha) which raise questions about the potential impact of technological interventions on farm income ([Harris and Orr, 2014](#)). Small farm sizes make farmers also more risk averse in trying out new technologies and practices. Second, there is a general lack of effective support for smallholder agriculture in much of sub-Saharan Africa, such that there are actually economic disincentives to investment in agriculture ([Ehui and Pender, 2005](#)). Private sector support is often uncertain, because only a small part of farm produce is marketed. Good markets for sale of extra produce and for input supply are a prerequisite condition for the widespread adoption of new agricultural technologies that aim at productivity increase. Third, failure of efficient extension services in most African countries limit the possibility to scale up best management practices ([Davis et al., 2008](#)). Fourth, effective policies, including input subsidies, which incentivize farmers to invest in new technologies and practices are needed ([Jayne and Rashid, 2013](#)).

Farming systems and farmers in sub-Saharan Africa being highly diverse, a major challenge with the development and dissemination of practices such as conservation agriculture and agroforestry is the identification of situations where they can offer major benefits ([Giller et al., 2011](#); [Luedeling et al., 2011](#)). Specific technologies should be adapted and targeted to categories of farmers who are likely to benefit from these technologies, and adapted to the local circumstances and specific farming contexts. Features that need to be considered include farmers' production objectives and constraints, expected costs (requirements in terms of inputs, equipment, labor), benefits (especially in the short term), production and financial risk, input supply and marketing, farmers' perceptions, and technical advice. This can be achieved through an innovation systems approach that fosters dynamic interactions and collaboration among researchers, extension agents, farmers, policy makers and the private sector ([Schut et al., 2016](#)). More generally, an alignment of agriculture and climate policy also appears to be essential ([Chabbi et al., 2017](#)).

4.4. Recommendations for SOC monitoring in field experiments

Implementing the 4‰ initiative on a large scale will require the capacity to measure and monitor SOC stocks with acceptable accuracy, quantifiable uncertainty and at relatively low cost ([Paustian et al., 2016](#)). This implies the combination of direct measurements with modeling approaches ([Post et al., 2001](#)). For example, repeated measurements at a given site are needed to establish relationships between environmental and management factors and SOC dynamics to improve statistical and quantitative models as tools in monitoring, reporting and verification systems for carbon in soils ([Batjes and van Wesemael, 2015](#)). Our review clearly highlighted the scarcity of good quality data on SOC stocks under agroforestry and conservation agriculture practices in sub-Saharan Africa.

In general, there is a lack of standardized procedures for studying and reporting SOC storage, making it difficult to evaluate the precise effect of a given management practice. We recommend the use of the diachronic approach to estimate SOC storage rates given the high uncertainty with the synchronic approach because of spatial variability of

soil properties that may influence SOC storage rates (e.g. Cerri et al., 2004). A prerequisite for interpretation of reported SOC data is information on the experimental design together with soil sampling methods (including sampling depth), sample size and a measure of variance of SOC stocks. Soil sampling depth was reported for different soil depths across the studies that we analyzed. Although the 2006 IPCC guidelines recommend a reference depth of 30 cm, it is useful to have estimates for SOC stocks of the entire root zone or to a depth of 1 or 2 m, especially in agroforestry systems (Cardinael et al., 2015a, 2018). Amongst other minimum data to be reported are the exact location of the experiment, climate, soil type and texture, soil bulk density, land-use history, date of establishment of the SOC-restoration practice, land management practices (e.g. fertilizer and organic residue applications), tree density and spacing as well as the management of pruning residues in the case of agroforestry systems. Finally, an estimate of the plant productivity in the experiments is helpful for interpretation of measured SOC storage rates as it determines the carbon inputs into soil.

5. Conclusions

Our systematic review of published literature showed that both agroforestry and conservation agriculture have a large potential to increase SOC stocks above the rate of $4\% \text{ yr}^{-1}$ in sub-Saharan Africa. Respectively 80 and 64% of the observations in the published studies had rates that were larger than $4\% \text{ yr}^{-1}$ (Fig. 2). The data showed, however, a very large variability. As a result, we found that only the agroforestry practice of fallows and multistrata systems led to SOC storage rates significantly higher than $4\% \text{ yr}^{-1}$ (Fig. 3). Among the agroforestry systems, parklands showed the lowest SOC storage rates, a median rate of $8\% \text{ yr}^{-1}$. Only 'full' conservation agriculture, i.e. with all three principles implemented, resulted in SOC storage rates that were significantly higher than $4\% \text{ yr}^{-1}$. The lowest rates were attained with no-tillage without residue retention.

The example of SOC gradients across different fields of individual farms and across farms illustrates the potential and challenges in increasing SOC stocks in the context of African smallholder agriculture. Low SOC levels are to a great extent the result of the limited resources of smallholder farmers in sub-Saharan Africa. Since climate change mitigation is not a priority for these smallholders, the 4% initiative has to be implemented on the grounds of the co-benefits that are offered through improved SOC storage. It means that for the African situation SOC-restoration practices should be promoted on the basis of their positive impact on crop productivity. Furthermore, these practices need to be targeted and adapted to the specific bio-physical and socio-economic situations of the farmers. The strategies to implement the 4% initiative in sub-Saharan Africa include the promotion of good markets for sale of extra produce and for input supply, effective private support and policy, including credit schemes and subsidies for inputs, and efficient extension services which incentivize farmers to invest in new technologies.

In conclusion, we argue that there is a great potential for SOC storage in agricultural soils of sub-Saharan Africa given the large areas of SOC-depleted soils. Practices such as agroforestry and conservation agriculture have a potential to restore SOC in these soils. The efficiency in doing so will depend on the specific situations and will need economic support to smallholder farmers. Climate change mitigation should be seen as a no-regret option, not as a key driver for promoting best management options. The key driver is increased farm income through higher crop productivity.

Acknowledgements

This work was partly funded by the French Agricultural Research Centre for International Development (CIRAD) and the CGIAR Research Program MAIZE, FP 4: Sustainable Intensification of Maize-based Cropping Systems. The authors are grateful to Philippe Letourmy for his

help with the statistical analysis.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.still.2018.02.015>.

References

- Abaker, W.E., Berninger, F., Saiz, G., Braojos, V., Starr, M., 2016. Contribution of *Acacia senegal* to biomass and soil carbon in plantations of varying age in Sudan. *For. Ecol. Manage.* 368, 71–80.
- Agbede, T.M., 2008. Nutrient availability and cocoyam yield under different tillage practices. *Soil Tillage Res.* 99, 49–57.
- Agbenin, J.O., Goladi, J.T., 1997. Carbon, nitrogen and phosphorus dynamics under continuous cultivation as influenced by farmyard manure and inorganic fertilizers in the savanna of northern Nigeria. *Agric. Ecosyst. Environ.* 63, 17–24.
- Agele, S.O., Ewulo, B.S., Oyewusi, I.K., 2005. Effects of some soil management systems on soil physical properties, microbial biomass and nutrient distribution under rainfed maize production in a humid rainforest Alfisol. *Nutr. Cycl. Agroecosyst.* 72, 121–134.
- Angers, D.A., Eriksen-Hamel, N.S., 2008. Full-inversion tillage and organic carbon distribution in soil profiles: a meta-analysis. *Soil Sci. Soc. Am. J.* 72, 1370.
- Anikwe, M.A.N., Ubochi, J.N., 2007. Short-term changes in soil properties under tillage systems and their effect on sweet potato (*Ipomea batatas* L.) growth and yield in an Ultisol in south-eastern Nigeria. *Aust. J. Soil Res.* 45 (5), 351–358.
- Baker, J.M., Ochsner, T.E., Venterea, R.T., Griffis, T.J., 2007. Tillage and soil carbon sequestration—what do we really know? *Agric. Ecosyst. Environ.* 118, 1–5.
- Balesdent, J., Balabane, M., 1996. Major contribution of roots to soil carbon storage inferred from maize cultivated soils. *Soil Biol. Biochem.* 28, 1261–1263.
- Barré, P., Angers, D.A., Basile-Doelsch, I., Bispo, A., Cécillon, L., Chenu, C., Chevallier, T., Derrien, D., Eglin, T.K., Pellerin, S., 2017. Ideas and perspectives: can we use the soil carbon saturation deficit to quantitatively assess the soil carbon storage potential, or should we explore other strategies? *Biogeosci. Discuss.* 1–12.
- Bationo, A., Kihara, J., Vanlauwe, B., Waswa, B., Kimetu, J., 2007. Soil organic carbon dynamics, functions and management in West African agro-ecosystems. *Agric. Syst.* 94, 13–25.
- Batjes, N.H., 1996. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.* 47, 151–163.
- Batjes, N.H., van Wesemael, B., 2015. Measuring and monitoring soil carbon. In: Banwart, S.A., Noellemeier, E., Milne, E. (Eds.), *Soil Carbon: Science, Management and Policy for Multiple Benefits*. CABI, Wallingford, UK, pp. 188–201.
- Baumert, S., Khamzina, A., Vlek, P.L.G., 2016. Soil organic carbon sequestration in *Jatropha curcas* Systems in Burkina Faso. *Land Degrad. Dev.* 27, 1813–1819.
- Blanco-Canqui, H., Lal, R., 2007. Soil structure and organic carbon relationships following 10 years of wheat straw management in no-till. *Soil Tillage Res.* 95, 240–254.
- Bright, M.B.H., Diedhiou, I., Bayala, R., Assigbetse, K., Chapuis-Lardy, L., Ndour, Y., Dick, R.P., 2017. Long-term *Piliostigma reticulatum* Intercropping in the sahel: crop productivity, carbon sequestration, nutrient cycling, and soil quality. *Agric. Ecosyst. Environ.* 242, 9–22.
- Cardinael, R., Chevallier, T., Barthès, B.G., Saby, N.P., Parent, T., Dupraz, C., Bernoux, M., Chenu, C., 2015a. Impact of alley cropping agroforestry on stocks, forms and spatial distribution of soil organic carbon - a case study in a mediterranean context. *Geoderma* 259–260, 288–299.
- Cardinael, R., Mao, Z., Prieto, I., Stokes, A., Dupraz, C., Kim, J.H., Jourdan, C., 2015b. Competition with winter crops induces deeper rooting of walnut trees in a mediterranean alley cropping agroforestry system. *Plant Soil* 391, 219–235.
- Cardinael, R., Guenet, B., Chevallier, T., Dupraz, C., Cozzi, T., Chenu, C., 2018. High organic inputs explain shallow and deep SOC storage in a long-term agroforestry system - combining experimental and modeling approaches. *Biogeosciences* 15, 297–317.
- Carter, S.E., Murwira, H.K., 1995. Spatial variability in soil fertility management and crop response in Mutoko Communal area, Zimbabwe. *Ambio* 24, 77–84.
- Castellano, M.J., Mueller, K.E., Olk, D.C., Sawyer, J.E., Six, J., 2015. Integrating plant litter quality, soil organic matter stabilization and the carbon saturation concept. *Global Change Biol.* 21, 3200–3209.
- Cerri, C.E.P., Cerri, C.C., Paustian, K., Bernoux, M., Mellilo, J.M., 2004. Combining soil C and N spatial variability and modeling approaches for measuring and monitoring soil carbon sequestration. *Environ. Manage.* 33, 274–288.
- Chabbi, A., Lehmann, J., Ciais, P., Loescher, H.W., Cotrufo, M.F., Don, A., Sancelme, M., Schipper, L., Six, J., Smith, P., Rumpel, C., 2017. Aligning agriculture and climate policy. *Nat. Clim. Change* 7, 307–309.
- Costa Junior, C., Corbeels, M., Bernoux, M., Piccolo, M.C., Neto, M.S., Feigl, B.J., Cerri, C.E.P., Cerri, C.C., Scopel, E., Lal, R., 2013. Assessing soil carbon storage rates under no-tillage: comparing the synchronic and diachronic approaches. *Soil Tillage Res.* 134, 207–212.
- Davis, K.E., Ekboir, J., Spielman, D.J., 2008. Strengthening agricultural education and training in sub-Saharan Africa from an innovation systems perspective: a case study of Mozambique. *J. Agric. Educ. Ext.* 14, 35–51.
- de Vries, W., 2018. Soil carbon 4 per mille: a good initiative but let's manage not only the soil but also the expectations. *Geoderma* 309, 111–112.
- de Vries, W., 2014. Nutrients trigger carbon storage. *Nat. Clim. Change* 4, 425–427.
- Dignac, M.-F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T., Freschet, G.T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron,

- P.-A., Nunan, N., Roumet, C., Basile-Doelsch, I., 2017. Increasing soil carbon storage: mechanisms, effects of agricultural practices and proxies. *A review. Agron. Sustain. Dev.* 37, 14.
- Ehui, S., Pender, J., 2005. Resource degradation, low agricultural productivity, and poverty in sub-Saharan Africa: pathways out of the spiral. *Agric. Econ.* 32, 225–242.
- FAO, 2015. *Conservation agriculture* [www document, URL <http://www.fao.org/ag/ca/1a.html>, (accessed 28/2/2018)].
- Feliciano, D., Ledo, A., Hillier, J., Nayak, D.R., 2018. Which agroforestry options give the greatest soil and above ground carbon benefits in different world regions? *Agric. Ecosyst. Environ.* 254, 117–129.
- Field, C.B., Mach, K.J., 2017. Rightsizing carbon dioxide removal. *Science* 356, 706–707.
- Galecki, A., Burzykowski, T., 2013. *Linear mixed-effects models using R: a step-by-step approach*. Springer Texts in Statistics.
- Gelaw, A.M., Singh, B.R., Lal, R., 2014. Soil organic carbon and total nitrogen stocks under different land uses in a semi-arid watershed in Tigray, Northern Ethiopia. *Agric. Ecosyst. Environ.* 188, 256–263.
- Giller, K.E., Tittonell, P., Rufino, M.C., van Wijk, M.T., Zingore, S., Mapfumo, P., Adjei-Nsiah, S., Herrero, M., Chikowo, R., Corbeels, M., Rowe, E.C., Baijuyka, F., Mwijage, A., Smith, J., Yeboah, E., van der Burg, W.J., Sanogo, O.M., Misiko, M., de Ridder, N., Karanja, S., Kaizzi, C., K'ungu, J., Mwale, M., Nwaga, D., Pacini, C., Vanlauwe, B., 2011. Communicating complexity: integrated assessment of trade-offs concerning soil fertility management within african farming systems to support innovation and development. *Agric. Syst.* 104, 191–203.
- Giller, K.E., Witter, E., Corbeels, M., Tittonell, P., 2009. Conservation agriculture and smallholder farming in Africa: the heretics' view. *Food Crops Res.* 114, 23–34.
- Gnankambary, Z., Bayala, J., Malmer, A., Nyberg, G., Hien, V., 2008. Decomposition and nutrient release from mixed plant litters of contrasting quality in an agroforestry parkland in the south-Sudanese zone of West Africa. *Nutr. Cycl. Agroecosyst.* 82, 1–13.
- Haile, S.G., Nair, V.D., Nair, P.K.R., 2010. Contribution of trees to carbon storage in soils of silvopastoral systems in Florida, USA. *Global Change Biol.* 16, 427–438.
- Harris, D., Orr, A., 2014. Is rainfed agriculture really a pathway from poverty? *Agric. Syst.* 123, 84–96.
- Hassink, J., 1997. The capacity of soils to preserve organic C and N by their association with clay and silt particles. *Plant Soil* 191, 77–87.
- Henry, M., Valentini, R., Bernoux, M., 2009. Soil carbon stocks in ecoregions of Africa. *Biogeosci. Discuss.* 6, 797–823.
- Hobbs, P.R., Govaerts, B., 2010. How conservation agriculture can contribute to buffering climate change. In: Reynolds, M.P. (Ed.), *Climate Change and Crop Production*. CAB International, pp. 177–199.
- Jayne, T.S., Rashid, S., 2013. Input subsidy programs in sub-Saharan Africa: a synthesis of recent evidence. *Agric. Econ. (United Kingdom)* 44, 547–562.
- Jerneck, A., Olsson, L., 2014. Food first! Theorising assets and actors in agroforestry: risk evaders, opportunity seekers and “the food imperative” in sub-Saharan Africa. *Int. J. Agric. Sustain.* 12, 1–22.
- Jerneck, A., Olsson, L., 2013. More than trees! Understanding the agroforestry adoption gap in subsistence agriculture: insights from narrative walks in Kenya. *J. Rural Stud.* 32, 114–125.
- Kimaro, A., Isaac, M., Chamshama, S., 2011. Carbon pools in tree biomass and soils under rotational woodlot systems in eastern Tanzania. In: Kumar, B., Nair, P. (Eds.), *Carbon Sequestration Potential of Agroforestry Systems*. Springer, pp. 142–156.
- Kintché, K., Guibert, H., Sogbedji, J.M., Levêque, J., Tittonell, P., 2010. Carbon losses and primary productivity decline in savannah soils under cotton-cereal rotations in semi-arid Togo. *Plant Soil* 336, 469–484.
- Kirkby, C.A., Richardson, A.E., Wade, L.J., Batten, G.D., Blanchard, C., Kirkegaard, J.A., 2013. Carbon-nutrient stoichiometry to increase soil carbon sequestration. *Soil Biol. Biochem.* 60, 77–86.
- Kirkby, C.A., Richardson, A.E., Wade, L.J., Passioura, J.B., Batten, G.D., Blanchard, C., Kirkegaard, J.A., 2014. Nutrient availability limits carbon sequestration in arable soils. *Soil Biol. Biochem.* 68, 402–409.
- Laganière, J., Angers, D.A., Paré, D., 2010. Carbon accumulation in agricultural soils after afforestation: a meta-analysis. *Global Change Biol.* 16, 439–453.
- Lal, R., 2016. Beyond COP 21: potential and challenges of the “4 per thousand” initiative. *J. Soil Water Conserv.* 71, 20A–25A.
- Lal, R., 2015. Sequestering carbon and increasing productivity by conservation agriculture. *J. Soil Water Conserv.* 70, 55A–62A.
- Lal, R., 2006. Enhancing crop yields in the developing countries through restoration of the soil organic carbon pool in agricultural lands. *Land Degrad. Dev.* 17, 197–209.
- Lal, R., 2004. Soil carbon sequestration impacts on global climate change and food security. *Science* 304, 1623–1627.
- Lal, R., 1998. Soil quality changes under continuous cropping for seventeen seasons of an Alfisol in western Nigeria. *Land Degrad. Dev.* 9, 259–274.
- Le Quéré, C., Andrew, R.M., Canadell, J.G., Sitch, S., Korsbakken, J.I., Peters, G.P., Manning, A.C., Boden, T.A., Tans, P.P., Houghton, R.A., Keeling, R.F., Alin, S., Andrews, O.D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L.P., Ciais, P., Currie, K., Delire, C., Doney, S.C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A.K., Kato, E., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozi, D., Melton, J.R., Metz, L., Miller, F., Monteiro, P.M.S., Munro, D.R., Nabel, J.E.M.S., Nakaoka, S., O'Brien, K., Olsen, A., Omar, A.M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B.D., Sutton, A.J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx, I.T., van der Werf, G.R., Viovy, N., Walker, A.P., Wiltshire, A.J., Zaehle, S., 2016. Global carbon budget 2016. *Earth Syst. Sci. Data* 8, 605–649.
- Le Quéré, C., Peters, G.P., Andres, R.J., Andrew, R.M., Boden, T., Ciais, P., Friedlingstein, P., Houghton, R.A., Marland, G., Moriarty, R., Sitch, S., Tans, P., Arneeth, A., Arvanitis, A., Bakker, D.C.E., Bopp, L., Canadell, J.G., Chini, L.P., Doney, S.C., Harper, A., Harris, I., House, J.I., Jain, A.K., Jones, S.D., Kato, E., Keeling, R.F., Klein Goldewijk, K., Körtzinger, A., Koven, C., Lefèvre, N., Omar, A., Ono, T., Park, G.-H., Pfeil, B., Poulter, B., Raupach, M.R., Regnier, P., Rödenbeck, C., Saito, S., Schwinger, J., Segsneider, J., Stocker, B.D., Tilbrook, B., van Heuven, S., Viovy, N., Wanninkhof, R., Wiltshire, A., Zaehle, S., Yue, C., 2015. Global carbon budget 2014. *Earth Syst. Sci. Data* 7, 45–85.
- Lemenih, M., Karlun, E., Olsson, M., 2005. Soil organic matter dynamics after deforestation along a farm field chronosequence in southern highlands of Ethiopia. *Agric. Ecosyst. Environ.* 109, 9–19.
- Lenth, R.V., 2016. Least-squares means: the R package lsmeans. *J. Stat. Softw.* 69, 1–33.
- Littell, R.C., Milliken, G.A., Walter, S.W., Wolfinger, R.D., Schabenberger, O., 2006. *SAS for Mixed Models*, second edition. SAS Institute.
- Lorenz, K., Lal, R., 2014. Soil organic carbon sequestration in agroforestry systems. A review. *Agron. Sustain. Dev.* 34, 443–454.
- Luedeling, E., Neufeldt, H., 2012. Carbon sequestration potential of parkland agroforestry in the Sahel. *Clim. Change* 115, 443–461.
- Luedeling, E., Sileshi, G., Beedy, T., Dietz, J., 2011. Carbon sequestration potential of agroforestry systems in Africa. In: Kumar, B.M., Nair, P.K.R. (Eds.), *Carbon Sequestration Potential of Agroforestry Systems*. Springer, pp. 61–83.
- Luo, Z., Wang, E., Sun, O.J., 2010. Can no-tillage stimulate carbon sequestration in agricultural soils? A meta-analysis of paired experiments. *Agric. Ecosyst. Environ.* 139, 224–231.
- Makumba, W., Akinnifesi, F.K., Janssen, B., Oenema, O., 2007. Long-term impact of a gliricidia-maize intercropping system on carbon sequestration in southern Malawi. *Agric. Ecosyst. Environ.* 118, 237–243.
- Meinshausen, M., Meinshausen, N., Hare, W., Raper, S.C.B., Frieler, K., Knutti, R., Frame, D.J., Allen, M.R., 2009. Greenhouse-gas emission targets for limiting global warming to 2°C. *Nature* 458, 1158–1162.
- Menichetti, L., Ekblad, A., Kätterer, T., 2015. Contribution of roots and amendments to soil carbon accumulation within the soil profile in a long-term field experiment in Sweden. *Agric. Ecosyst. Environ.* 200, 79–87.
- Minasny, B., Malone, B.P., McBratney, A.B., Angers, D.A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B.S., Field, D.J., Gimona, A., Hedley, C.B., Hong, S.Y., Mandal, B., Marchant, B.P., Martin, M., McConkey, B.G., Mulder, V.L., O'Rourke, S., Richer-de-Forges, A.C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovov, V., Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., van Wesemael, B., Winowiecki, L., 2017. Soil carbon 4 per mille. *Geoderma* 292, 59–86.
- Mitchell, R.J., Campbell, C.D., Chapman, S.J., Cameron, C.M., 2010. The ecological engineering impact of a single tree species on the soil microbial community. *J. Ecol.* 98, 50–61.
- Mloza-Banda, M.L., Cornelis, W.M., Mloza-Banda, H.R., Makwiza, C.N., Verbist, K., 2014. Soil properties after change to conservation agriculture from ridge tillage in sandy clay loams of mid-altitude Central Malawi. *Soil Use Manage.* 30, 569–578.
- Moebius-Clune, B.N., van Es, H.M., Idowu, O.J., Schindelbeck, R.R., Kimetu, J.M., Ngoze, S., Lehmann, J., Kinyangi, J.M., 2011. Long-term soil quality degradation along a cultivation chronosequence in western Kenya. *Agric. Ecosyst. Environ.* 141, 86–99.
- Nair, P.K.R., 1993. *An Introduction to Agroforestry*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Nair, P.K.R., 1985. Classification of agroforestry systems. *Agrofor. Syst.* 3, 97–128.
- Nair, P.K.R., Buresh, R., Mugendi, D.N., Latt, C.R., 1999. Nutrient cycling in tropical agroforestry systems: myths and science. In: Buck, L.E., Lassoie, J.P., Fernandes, E.C.M. (Eds.), *Agroforestry in Sustainable Agricultural Systems*. CRC Press LLC, pp. 1–32.
- Nair, P.K.R., Nair, V.D., 2014. Solid-fluid-gas: the state of knowledge on carbon-sequestration potential of agroforestry systems in Africa. *Curr. Opin. Environ. Sustain.* 6, 22–27.
- Ngome, A.F., Becker, M., Mtei, K.M., Musngungu, F., 2011. Fertility management for maize cultivation in some soils of Western Kenya. *Soil Tillage Res.* 117, 69–75.
- Nyamadzawo, G., Chikowo, R., Nyamugafata, P., Giller, K.E., 2007. Improved legume tree fallows and tillage effects on structural stability and infiltration rates of a kaolinitic sandy soil from central Zimbabwe. *Soil Tillage Res.* 96, 182–194.
- Nyamadzawo, G., Chikowo, R., Nyamugafata, P., Nyamangara, J., Giller, K.E., 2008. Soil organic carbon dynamics of improved fallow-maize rotation systems under conventional or no-tillage in Central Zimbabwe. *Nutr. Cycl. Agroecosyst.* 81, 85–93.
- Ojeniyi, S.O., Adekayode, F.O., 1999. Soil conditions and cowpea and maize yield produced by tillage methods in the rainforest zone of Nigeria. *Soil Tillage Res.* 51, 161–164.
- Paul, B.K., Vanlauwe, B., Hoogmoed, M., Hurisso, T.T., Ndamanyenye, T., Terano, Y., Six, J., Ayuke, F.O., Pulleman, M.M., 2015. Exclusion of soil macrofauna did not affect soil quality but increased crop yields in a sub-humid tropical maize-based system. *Agric. Ecosyst. Environ.* 208, 75–85.
- Paustian, K., Collins, H.P., Paul, E.A., 1997. Management controls on soil carbon. In: Paul, E.A., Paustian, K., Elliott, E.T., Cole, C.V. (Eds.), *Soil organic matter in temperate agroecosystems*. CRC Press, Boca Raton, pp. 15–49.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G.P., Smith, P., 2016. Climate-smart soils. *Nature* 532, 49–57.
- Peichl, M., Thevathasan, N.V., Gordon, A.M., Huss, J., Abohassan, R.A., 2006. Carbon sequestration potentials in temperate tree-based intercropping systems, southern Ontario, Canada. *Agrofor. Syst.* 66, 243–257.
- Pierret, A., Maeght, J.-L., Clément, C., Montoro, J.-P., Hartmann, C., Gonkhamdee, S., 2016. Understanding deep roots and their functions in ecosystems: an advocacy for more unconventional research. *Ann. Bot.* 118, 621–635.
- Post, W.M., Izaurralde, R.C., Mann, L.K., Bliss, N., 2001. Monitoring and verifying changes of organic carbon in soil. *Clim. Change* 51, 73–99.

- Powelson, D.S., Stirling, C.M., Thierfelder, C., White, R.P., Jat, M.L., 2016. Does conservation agriculture deliver climate change mitigation through soil carbon sequestration in tropical agro-ecosystems? *Agric. Ecosyst. Environ.* 220, 164–174.
- Powelson, D.S., Whitmore, A.P., Goulding, K.W.T., 2011. Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *Eur. J. Soil Sci.* 62, 42–55.
- Prudencio, C., 1993. Ring management of soils and crops in the west African semi-arid tropics: the case of the mossi farming system in Burkina Faso. *Agric. Ecosyst. Environ.* 46, 1–17.
- Raddad, E.Y., Luukkanen, O., Salih, A.A., Kaarakka, V., Elfadl, M.A., 2006. Productivity and nutrient cycling in young *Acacia senegal* farming systems on vertisol in the Blue Nile region. *Sudan. Agrofor. Syst.* 68, 193–207.
- R Development Core Team, 2013. R: A Language and Environment for Statistical Computing.
- Ranaivoson, L., Naudin, K., Ripoche, A., Affholder, F., Rabeharisoa, L., Corbeels, M., 2017. Agro-ecological functions of crop residues under conservation agriculture. *Agron. Sustain. Dev.* 37 (26), 1–17.
- Richardson, A.E., Kirkby, C.A., Banerjee, S., Kirkegaard, J.A., 2014. The inorganic nutrient cost of building soil carbon. *Carbon Manage.* 5, 265–268.
- Rimhanen, K., Kotoja, E., Yli-Halla, M., Kahiluoto, H., 2016. Ethiopian agriculture has greater potential for carbon sequestration than previously estimated. *Global Change Biol.* 22, 3739–3749.
- Scharlemann, J.P., Tanner, E.V., Hiederer, R., Kapos, V., 2014. Global soil carbon: understanding and managing the largest terrestrial carbon pool. *Carbon Manage.* 5, 81–91.
- Scheffé, H., 1959. *The Analysis of Variance*. John Wiley & Sons, Inc.
- Schlesinger, W.H., 1977. Carbon balance in terrestrial detritus. *Annu. Rev. Ecol. Syst.* 8, 51–81.
- Schut, M., van Asten, P., Okafor, C., Hicintuka, C., Mapatano, S., Nabahungu, N.L., Kagabo, D., Muchunguzi, P., Njukwe, E., Dontop-Nguezet, P.M., Sartas, M., Vanlauwe, B., 2016. Sustainable intensification of agricultural systems in the Central African highlands: the need for institutional innovation. *Agric. Syst.* 145, 165–176.
- Sleshi, G., Akinnifesi, F.K., Ajayi, O.C., Place, F., 2008. Meta-analysis of maize yield response to woody and herbaceous legumes in sub-Saharan Africa. *Plant Soil* 307, 1–19.
- Six, J., Conant, R.T., Paul, E.A., Paustian, K., 2002. Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant Soil* 241, 155–176.
- Six, J., Elliott, E.T., Paustian, K., 1999. Aggregate and soil organic matter dynamics under conventional and no-tillage systems. *Soil Sci.* 63, 1350–1358.
- Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, B., Sirotenko, O., Howden, M., McAllister, T., Pan, G., Romanenkov, V., Schneider, U., Towprayoon, S., Wattenbach, M., Smith, J., 2008. Greenhouse gas mitigation in agriculture. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 363, 789–813.
- Solomon, D., Fritzsche, F., Lehmann, J., Tekalign, M., Zech, W., 2002. Soil organic matter dynamics in the subhumid agroecosystems of the Ethiopian highlands: evidence from natural C-13 abundance and particle-size fractionation. *Soil Sci. Soc. Am. J.* 66, 969–978.
- Solomon, D., Lehmann, J., Kinyangi, J., Amelung, W., Lobe, I., Pell, A., Riha, S., Ngoze, S., Verchot, L., Mbugua, D., Skjemstad, J., Schäfer, T., 2007. Long-term impacts of anthropogenic perturbations on dynamics and speciation of organic carbon in tropical forest and subtropical grassland ecosystems. *Global Change Biol.* 13, 511–530.
- Stewart, C.E., Paustian, K., Conant, R.T., Plante, A.F., Six, J., 2007. Soil carbon saturation: concept, evidence and evaluation. *Biogeochemistry* 86, 19–31.
- Stockmann, U., Adams, M.A., Crawford, J.W., Field, D.J., Henakaarchchi, N., Jenkins, M., Minasy, B., McBratney, A.B., Courcelles, V.D.R.D., Singh, K., Wheeler, I., Abbott, L., Angers, D.A., Baldock, J., Bird, M., Brookes, P.C., Chenu, C., Jastrow, J.D., Lal, R., Lehmann, J., O'Donnell, A.G., Parton, W.J., Whitehead, D., Zimmermann, M., 2013. The knowns, known unknowns and unknowns of sequestration of soil organic carbon. *Agric. Ecosyst. Environ.* 164, 80–99.
- Swanepoel, C.M., van der Laan, M., Weepener, H.L., du Preez, C.C., Annandale, J.G., 2016. Review and meta-analysis of organic matter in cultivated soils in southern Africa. *Nutr. Cycl. Agroecosyst.* 104, 107–123.
- Takimoto, A., Nair, P.K.R., Nair, V.D., 2008. Carbon stock and sequestration potential of traditional and improved agroforestry systems in the West African Sahel. *Agric. Ecosyst. Environ.* 125, 159–166.
- Tittonell, P., Muriuki, A., Klapwijk, C.J., Shepherd, K.D., Coe, R., Vanlauwe, B., 2013. Soil heterogeneity and soil fertility gradients in smallholder farms of the East African highlands. *Soil Sci. Soc. Am. J.* 77, 525–538.
- Tittonell, P., Vanlauwe, B., Corbeels, M., Giller, K.E., 2008. Yield gaps, nutrient use efficiencies and response to fertilisers by maize across heterogeneous smallholder farms of western Kenya. *Plant Soil* 313, 19–37.
- Tittonell, P., Vanlauwe, B., Leffelaar, P.A., Shepherd, K.D., Giller, K.E., 2005. Exploring diversity in soil fertility management of smallholder farms in western Kenya. *Agric. Ecosyst. Environ.* 110, 166–184.
- Torn, M.S., Trumbore, S.E., Chadwick, O.A., Vitousek, P.M., Hendricks, D.M., 1997. Mineral control of soil organic carbon storage and turnover. *Nature* 389, 3601–3603.
- Torquebiau, E.F., 2000. A renewed perspective on agroforestry concepts and classification. *Life Sci.* 323, 1009–1017.
- Tumwebaze, S.B., Bevilacqua, E., Briggs, R., Volk, T., 2011. Soil organic carbon under a linear simultaneous agroforestry system in Uganda. *Agrofor. Syst.* 84, 11–23.
- Vågen, T.-G., Lal, R., Singh, B.R., 2005. Soil carbon sequestration in sub-Saharan Africa: a review. *Land Degrad. Dev.* 16, 53–71.
- van Groenigen, J.W., van Kessel, C., Hungate, B.A., Oenema, O., Powelson, D.S., van Groenigen, K.J., 2017. Sequestering soil organic carbon: a nitrogen dilemma. *Environ. Sci. Technol.* 51, 4738–4739.
- van Groenigen, K.-J., Six, J., Hungate, B.A., de Graaff, M.-A., van Breemen, N., van Kessel, C., 2006. Element interactions limit soil carbon storage. *PNAS* 103, 6571–6574.
- VandenBygaert, A.J., 2018. Comments on soil carbon 4 per mille by Minasy et al. 2017. *Geoderma* 309, 113–114.
- VandenBygaert, A.J., Kay, B.D., 2004. Persistence of soil organic carbon after plowing a long-term no-till field in southern Ontario, Canada. *Soil Sci. Soc. Am. J.* 68, 1394–1402.
- Vanlauwe, B., Diels, J., Sanginga, N., Merckx, R., 2005. Long-term integrated soil fertility management in South-western Nigeria: crop performance and impact on the soil fertility status. *Plant Soil* 273, 337–354.
- Verchot, L.V., Dutaour, L., Shepherd, K.D., Albrecht, A., 2011. Organic matter stabilization in soil aggregates: understanding the biogeochemical mechanisms that determine the fate of carbon inputs in soils. *Geoderma* 161, 182–193.
- Virto, I., Barré, P., Burlot, A., Chenu, C., 2012. Carbon input differences as the main factor explaining the variability in soil organic C storage in no-tilled compared to inversion tilled agrosystems. *Biogeochemistry* 108, 17–26.
- Wang, Z., Hoffmann, T., Six, J., Kaplan, J.O., Govers, G., Doetterl, S., Van Oost, K., 2017. Human-induced erosion has offset one-third of carbon emissions from land cover change. *Nat. Clim. Change* 1–6.
- Zingore, S., Manyame, C., Nyamugafata, P., Giller, K.E., 2005. Long-term changes in organic matter of woodland soils cleared for arable cropping in Zimbabwe. *Eur. J. Soil Sci.* 56, 727–736.
- Zingore, S., Murwira, H.K., Delve, R.J., Giller, K.E., 2007. Soil type, management history and current resource allocation: three dimensions regulating variability in crop productivity on African smallholder farms. *Food Crops Res.* 101, 296–305.
- Zingore, S., Tittonell, P., Corbeels, M., van Wijk, M.T., Giller, K.E., 2011. Managing soil fertility diversity to enhance resource use efficiencies in smallholder farming systems: a case from Murewa District, Zimbabwe. *Nutr. Cycl. Agroecosyst.* 90, 87–103.
- Zomer, R.J., Neufeldt, H., Xu, J., Ahrends, A., Bossio, D., Trabucco, A., van Noordwijk, M., Wang, M., 2016. Global tree cover and biomass carbon on agricultural Land: the contribution of agroforestry to global and national carbon budgets. *Sci. Rep.* 6, 29987. <http://dx.doi.org/10.1038/srep29987>.